

List of Figures and Tables

FIGURES

1.1	The main groupings of scientific specialists working on the material excavated from Çatalhöyük.	8
1.2	Overlaps between the research interests of the different funders at Çatalhöyük.	9
1.3	Specialist groups and their research networks.	9
2.1	Database infrastructure.	14
2.2	Diagram showing the defragmented recording model.	15
2.3	The Çatalhöyük Living Archive web application.	20
2.4	Multimodal search and browse in a spatial-temporal browser, to reconstitute a burial.	21
2.5	Searching both the RDF store and traditional relational database.	21
3.1	Social Network of the Çatalhöyük Team, 1993–2013.	27
3.2	Sample of topics modelled.	27
3.3	Document and Topic Network.	28
3.4	Network in 1994, illustrating the small size of the team and a disconnected social structure.	29
3.5	Network in 1996, illustrating the growth of the project team but few opportunities for information flow.	29
3.6	Network in 1998, illustrating the increased integration of the team.	30
3.7	Network in 1999, illustrating the effects of the six-month field season.	31
3.8	Network in 2000, illustrating the disaggregation associated with this and other study seasons.	32
3.9	Network in 2002, illustrating renewed cohesion in the team's social structure.	33
3.10	Network in 2003, illustrating the network breaking apart and new forms of collaboration emerging.	34
3.11	Network in 2009, illustrating the disintegration of the network during a study season.	36
4.1	Data capturing sessions via laser scanning and image-based 3D modelling.	44
4.2	Virtual excavation of Space 77, Feature 3686 (sk.20430).	45
4.3	3D surface model of B.89 generated in Agisoft Photoscan and implemented in the 3D GIS using GCPs to georeference the model.	47
4.4	3D GIS visualization of Mellaart phases superimposed to the models generated by IBM by the 3D-Digging Project.	49
4.5	Diverse datasets—acquired in different field campaigns—were implemented and visualized into the 3D GIS platform (ArcScene) during season 2013.	50
4.6	Immersive simulation of B.89 in the DiVE.	50
4.7	Ortho view (a) and perspective view (b) of B.89 in the 3D GIS of the South Area.	51
4.8	Observable data of unit19807.	52
4.9	X-ray shader applied to the 3D model of unit19807.	52
4.10	Main activities and affordances in the spatial domain of B.89.	53
4.11	3D print of human mandible 19829.X2 retrieved in B.89 in 2012.	53
4.12	Orthophoto of B.97 south wall section generated using image-based 3D Modelling.	54
4.13	Aligned point clouds of B.77 scanned in 2012.	55
5.1	Osteoarthritis of the knee joint as indicated by the presence of marginal lipping and fine porosity on the articular surface of the right and left patellae.	60
5.2	Medial epicondylitis of the right humerus as indicated by the presence of surface porosity and enthesophytes at the common flexor origin.	61
5.3	Example of a grinder from the Çatalhöyük assemblage.	62
5.4	Example of a grinding slab/quern from the Çatalhöyük assemblage.	62
5.5	Examples of projectile points from the Çatalhöyük assemblage.	63
5.6	Frequency and severity of hip osteoarthritis between males and females at Çatalhöyük.	65
5.7	Frequency and severity of ankle osteoarthritis between males and females at Çatalhöyük.	65
5.8	Frequency and severity of foot osteoarthritis between males and females at Çatalhöyük.	66
5.9	Frequency and severity of hand osteoarthritis between males and females at Çatalhöyük.	66
5.10	Ratio of lateral to medial epicondylitis (L/M) in the right and left arms of males and females at Çatalhöyük.	67
5.11	Weight distribution of complete grinders (n=29) during Period 1 and Period 2 at Çatalhöyük.	68
5.12	Size distribution of complete grinders (n=31) during Period 1 and Period 2 at Çatalhöyük.	68
5.13	Size distribution of complete grinding slabs/querns (n=23) during Period 1 and Period 2 at Çatalhöyük.	69
5.14	Distribution of arrowheads and spearheads between Period 1 and Period 2 at Çatalhöyük based on projectile point analysis using the Hildebrandt and King method (2012).	69
5.15	Ratio of lateral to medial epicondylitis (L/M) in the right and left arms of individuals dating to Period 1 or Period 2 at Çatalhöyük.	70
6.1	Assemblage of figurines showing emphasized buttocks, drooping breasts, and stomachs.	76
6.2	Human isotope data according to age stage.	79
6.3	Figuring 12401.X7, showing a fleshed front and skeletonized back.	81
6.4	Skeleton 10829 and 10813 with associated finds.	83
7.1	Mean isotope ratios for carbon and nitrogen indicate that diets between the sexes were essentially the same.	88
7.2	Adult tooth from Çatalhöyük individual showing evidence of caries.	89
7.3	A discrete patch of periosteal reactive bone indicative of non-specific infection on the right femur of an infant from Çatalhöyük.	89
7.4	Multiple healed rib fractures observed in a middle adult female (8115) from Neolithic Çatalhöyük.	89

7.5	Thin section of cortical bone in the rib of an adult individual from Çatalhöyük used to measure the amount and turnover of cortical bone tissue using histomorphometry.	90
7.6	% cortical bone (an indicator of the amount of bone cortex) in the rib across three broad age groups in the adults at Çatalhöyük.	90
7.7	Mean annual activation frequency (an indicator of bone turnover) in the rib across three broad age groups in the adults at Çatalhöyük.	91
7.8	Metacarpal cortical index across three broad age groups in the adults at Çatalhöyük.	91
7.9	The early focus by Mellaart and others on mother goddess imagery was largely based on the visual emphasis on figurines at Çatalhöyük, such as this well-known figurine of a seated female figure.	92
7.10	Female burial with fetus in situ excavated at Çatalhöyük.	92
8.1a	Obsidian bifaces/biface preform hoard (4209) in B.9, South H.	98
8.1b	Obsidian blade/spearhead preform hoard (1461) from B.1, North G	99
8.2	Refitting thinning flakes and Göllü Dağ biface (13111.x3) from B.60, North H.	100
8.3a	Thinning/retouch flakes and projectile fragments from point manufacture; Phase 1, B.56, South R.	101
8.3b	Thinning/retouch flakes and projectile fragments from point manufacture; Phase 1, B.56, South R.	101
8.4	(a) Reconstruction of a woman (11306) holding a plastered skull painted with ochre (11330); (b) the woman (11306) and plastered skull (11330) in situ.	102
8.5	Collection of material placed within the foundation of B.44, subsequently becoming the southwest platform.	103
8.6	Group of ground stone, worked bone, and obsidian (12807).	104
8.7	Plan of Sp.333, South P.	105
9.1	Overall plan of Çatalhöyük East Mound showing locations of the case study sequences.	112
9.2	Simplified model of the South Area sequence.	112
9.3	Microscopic components of floor within Sp.470.	114
9.4	Scheme of static and dynamic loads that may cause damage and deformation of architectural features.	115
9.5	Close-up view of Sp.470 and location of the related special deposits.	115
9.6	Simplified model of the North Area sequence.	116
9.7	Close-up view of Sp.488 and location of the related special deposits.	117
9.8	Close-up view of Sp.511 and location of the related special deposit and the collapsed remains.	118
9.9	Microscopic components of floor sequences present in the collapsed materials of Sp.511.	119
9.10	Microscopic features of ashy layers towards the top of roof/upper storey sequence	119
9.11	Simplified model of the TP Area sequence.	120
9.12	Close-up view of Sp.325/Sp.326 and the related special deposits.	122
10.1	Çatalhöyük site plan, showing the areas of study.	129
10.2	B.77, situated within the North Area at Çatalhöyük.	130
10.3	Plan of B.77 in its final phase, showing the bins and architecture as well as some of the rich artefact assemblages deposited prior to its final destruction by fire.	131
10.4	Overview of B.77 (south facing photograph).	131
10.5	Ochre hand prints on the north wall of B.77 (north facing photograph).	131
10.6	Bucrania and horned bench associated with the northeast platform of B.77 (northeast facing photograph).	132
10.7	In situ clusters of 'bone and stone' on the latest burnt floors of B.77 (southwest facing photograph).	132
10.8	Well-preserved bin structures surviving to the east of B.77 (north facing photograph).	132
10.9	Çatalhöyük Research Project database and intra-site GIS.	134
10.10	Schematic matrix with phase lines (red) and stratigraphic correlations.	135
10.11	Step 1—Vertical compression of the matrix.	135
10.12	Step 2—Calibration of the matrix by stratigraphic correlation.	135
10.13	Step 3—Final stratigraphic parse to establish unit lifespan.	136
10.14a	Animation 1—Basic sequence of animation stills visualizing the B.77 depositional sequence.	136
10.14b	Sequential frames of basic animated visualization of B.77 depositional sequence.	137
10.15a	Single still from the animation sequence visualizing the B.77 sequence and symbolized with basic depositional classification.	138
10.15b	Animation 2—Sequence of animation stills visualizing the B.77 sequence symbolized with basic depositional classifications.	139
10.16a	Single still from animation sequence visualizing B.77 and showing the integration of material culture types.	140
10.16b	Animation 3—Sequence of animation stills visualizing B.77 and showing the integration of material culture types.	141
10.17a	Single still from animation sequence visualizing B.77 and integrating preliminary statistical observations.	143
10.17b	Animation 4—Sequence of animation stills visualizing B.77 and integrating preliminary statistical observations.	144
10.18	Example of a frame from a previous case study animation (of the B.65 and B.56 sequences), which in this case shows the density of obsidian objects aligned adjacent to a graph of the same data.	145
10.19a	Single still from animation sequence visualizing B.77 and demonstrating a more complex integration of multiple datasets.	146
10.19b	Animation 5—Sequence of animation stills visualizing B.77 and demonstrating a more complex integration of multiple datasets.	147
11.1	TP Area and other excavation areas at Çatalhöyük East.	152
11.2	B.81 in the TP Area.	152
11.3	Clay use. Matching raw materials and the landscape.	153
11.4	Clay use at Çatalhöyük. Matching artefacts and raw materials.	154
11.5	Diagram of sheep second mandibular molar showing how sequential enamel sampling can provide a 12-month time capsule of isotopic data.	155
11.6	Sheep tooth occlusal surface (×8 resolution) showing area of dental microwear studied with examples of diet-generated striations and pits (×500 resolution).	155

11.7	Oxygen isotope curves constructed from sequential samples taken from second mandibular molars of modern sheep born in March and in May.	156
11.8	Modelling the Çatalhöyük landscape topography in the Early Neolithic.	157
11.9	Modelling the Çatalhöyük landscape topography in the Late Neolithic.	157
11.10	Vegetation zones in the Konya Plain.	158
11.11	Frequency of charcoal values of different wood species in the Early Neolithic.	159
11.12	Frequency of charcoal values of different wood species at the end of the Early and in the Late Neolithic.	160
11.13	Chart showing temporal trends in the birth month of TP and South Area sheep, based on modelled oxygen isotope evidence.	161
11.14	Chart showing temporal trends in the first year movement of TP and South Area sheep, based on modelled oxygen isotope evidence.	161
11.15	Chart showing temporal trends in the final dietary regime of TP and South Area sheep, based on modelled dental microwear evidence.	162
11.16	The incised panel above burial 327 in TP Area.	162
12.1	Femoral midshaft A–P/M–L bending strength (mean Zx/Zy) in males and females at Çatalhöyük and comparative Pleistocene and Holocene European samples.	169
12.2	Femoral midshaft A–P/M–L bending strength (mean Zx/Zy) in males and females across the three time periods of Çatalhöyük's occupation.	170
13.1	Map of excavation areas on the East Mound at Çatalhöyük.	180
13.2	Late Neolithic sites in central, western, and west-northern Anatolia.	181
13.3	Examples of Dark Gritty Ware and Light Local Ware from Mellaart's Levels III–II.	183
13.4	A typical holemouth jar from Mellaart's Level III.	184
13.5	Red slipped bowl with basket handle and relief from Mellaart's Level II, red painted sherd from TP P and base fragments from Levels III–II.	185
13.6	S-profiled developed bowls from KOPAL Area.	185
13.7	Examples of bowls and jars.	186
13.8	Examples of horizontally and vertically perforated lugs from Mellaart Levels III–II.	187
13.9	Examples of incised decoration from TP. 1–2—TP Q ₃ —TP M.	187
13.10	Selection of obsidian pressure blades and other implements from Late Neolithic Çatalhöyük.	189
13.11	Obsidian sources represented in the Late Neolithic chipped stone assemblage of Çatalhöyük.	190
13.12	Selection of obsidian projectiles and a retouched chert blade from Late Neolithic Çatalhöyük.	190

TABLES

4.1	Terrestrial laser scanning workflow at Çatalhöyük.	48
5.1	Scoring system for frequency and severity of osteoarthritis.	60
5.2	Comparison of the spear and the bow and arrow as hunting weapons.	64
5.3	Age-controlled frequency of osteoarthritis (% joints affected) for females and males at Çatalhöyük.	64
5.4	Ratio of osteoarthritis (% joints affected) in the right and left upper limbs of females and males as a measure of laterality.	66
5.5	Levels corresponding to the two time periods used in this analysis.	68
5.6	Age-controlled frequency of osteoarthritis (% joints affected) for Period 1 and Period 2 at Çatalhöyük.	70
8.1	Contextual information for a selection of obsidian hoards from the 1995–2008 seasons.	98
8.2	Foundation burials from Çatalhöyük by stratigraphic level.	102
11.1	The modelled use of oxygen isotope values in sheep teeth in identifying herding location during the first year of life.	156
11.2	Modelled use of dental microwear analysis in the interpretation of archaeological domestic sheep diets just before death.	157
11.3	Uses of clay for mudbrick, plaster, and pottery production at Çatalhöyük.	157
11.4	Summary of dental isotope and microwear results.	160
11.5	Two models of herd resource requirements associated with the breeding cycle, product goals, and labour demands	161
12.1	Levels corresponding to the three time periods used in this analysis.	170
12.2	Inferring mobility through the femoral midshaft index.	170